



Cannaroo

5000 Linbar Drive Suite 250
Nashville
TN, USA 37211
615-994-0660
carol@cannaroo.com



SAMPLE:D809092-04

METRC/Biotrack#N/A **Harvest/Lot ID: 1821912.8**
Batch#: 1821912.8, Batch Size: N/A -grams
Ordered: 9/23/2018 Sampled:9/23/2018
Completed: 9/25/2018 8:14:34 PM Expires: 09/25/19
Sampling Method: SOP SOP Client Method

Image



Safety


- Pesticides - NT
- Microbials - NT
- Mycotoxins - NT
- Heavy Metals - NT
- Terpenes - NT
- Residual-Solvents - NT
- Filth - NT
- Water Activity - NT
- Moisture - NT

Cannabinoids

Analyte	Weight(%)	mg/g
D9-THC	0.17	1.69
THCa	0.01	0.10
TOTAL THC	0.17	1.70
CBD	8.39	83.88
CBDa	0.01	0.12
TOTAL CBD	8.40	84.00
CBN	0.03	0.29
CBDV	0.08	0.79
D8-THC	0.01	0.10
THCV	ND	ND
CBG	0.05	0.50
CBGA	ND	ND
CBC	0.11	1.10
TOTAL CANNABINOIDS	8.86	88.57

Cannabinoids

0.17% Total THC	8.40% Total CBD
---------------------------	---------------------------





Cannaroo

5000 Linbar Drive Suite 250
Nashville
TN, USA 37211
615-994-0660
carol@cannaroo.com



SAMPLE:D809034-02

METRC/Biotrack#N/A **Harvest/Lot ID: 1821912.80**
Batch#: 1821912.80, Batch Size: N/A -grams
Ordered: 9/11/2018 Sampled:9/11/2018
Completed: 9/13/2018 7:14:14 PM Expires: 09/13/19
Sampling Method: SOP SOP Client Method

Cannabinoid Profile Test Result-Analysis Method :SOP.T.40.020, SOP.T.30.050

Reagent LOT ID #

Dilution id factor#C

Analytical Batch:B810055

filter id factor#C

Full spectrum cannabinoid analysis utilizing High Performance Liquid Chromatography with UV detection (HPLC-UV). (Method: SOP.T.30.050 for sample prep and Shimadzu High Sensitivity Method SOP.T.40.020 for analysis. LOQ for all cannabinoids is 1 mg/L).

Micro Analysis-Analysis method :SOP.T.40.043

Analytical Batch: B810053

Pathogens#

Aspergillus flavus
Aspergillus fumigatus
Aspergillus niger
Aspergillus terreus
Escherichia coli
Salmonella

Results#C

Absent in 1 gram
Absent in 1 gram
Absent in 1 gram
Absent in 1 gram
Absent in 1 gram
Absent in 1 gram

Microbiological testing for Fungal and Bacterial Identification via Polymerase Chain Reaction (PCR) method consisting of sample DNA amplified via tandem Polymerase Chain Reaction (PCR) as a crude lysate which avoids purification. (Method SOP.T.40.043) If a pathogenic Escherichia Coli, Salmonella, Aspergillus fumigatus, Aspergillus flavus, Aspergillus niger, or Aspergillus terreus is detected in 1g of a sample, the sample fails the microbiological-impurity testing.